

Applicant: Edward R. Burns and Vadiraja Murthy
Serial No.: 08/746,635
Filed: November 13, 1996
Page 2 of 5

Amendments to the Claims:

Please amend Claims 33, 34, and 35 as set forth below.

1-19. (Canceled)

20. (Previously presented) A method for diagnosing erythrocyte hemolysis in a subject who is suspected of having erythrocyte hemolysis *in vivo*, the method consisting essentially of the steps of:

(a) obtaining a serum sample from a subject who is suspected of having erythrocyte hemolysis *in vivo*; and

(b) determining the level of erythrocyte adenylate kinase activity in said sample, the presence of at least about 20 U/L erythrocyte adenylate kinase activity in said sample being indicative of erythrocyte hemolysis *in vivo* in said subject.

21-23. (Canceled)

24. (Previously presented) A method for diagnosing erythrocyte hemolysis in a subject who may have erythrocyte hemolysis, the method comprising the steps of:

(a) obtaining a serum sample from the subject; and

(b) determining the level of erythrocyte adenylate kinase activity in the sample by determining a value for total adenylate kinase activity in the serum sample;

determining a value for percentage erythrocyte adenylate kinase in the sample;

and

determining erythrocyte adenylate kinase activity in the sample by multiplying the

Applicant: Edward R. Burns and Vadiraja Murthy
Serial No.: 08/746,635
Filed: November 13, 1996
Page 3 of 5

value for percentage erythrocyte adenylate kinase by the value for total adenylate kinase activity,

the presence of at least about 20 U/L erythrocyte adenylate kinase activity in the sample being indicative of erythrocyte hemolysis in the subject.

25. (Canceled)

26. (Previously presented) A method for diagnosing erythrocyte hemolysis in a subject who may have erythrocyte hemolysis, the method comprising the steps of:

- (a) obtaining a serum sample from the subject; and
- (b) determining the level of erythrocyte adenylate kinase activity in the sample using an antibody that is specific for and binds to erythrocyte adenylate kinase, the presence of at least about 20 U/L erythrocyte adenylate kinase activity in the sample being indicative of erythrocyte hemolysis in the subject.

27-32. (Canceled)

33. (Currently amended) The method of claim 26, ~~where the~~ wherein erythrocyte adenylate kinase-bound antibody is detected using a radioimmunoassay.

34. (Currently amended) The method of claim 26, ~~where the~~ wherein erythrocyte adenylate kinase-bound antibody is detected using an enzyme-linked immunosorbent assay (ELISA).

Applicant: Edward R. Burns and Vadiraja Murthy

Serial No.: 08/746,635

Filed: November 13, 1996

Page 4 of 5

35. (Currently amended) The method of claim 26, ~~where the~~ wherein
erythrocyte adenylate kinase-bound antibody is detected using immunoprecipitation.

36. (Previously presented) The method of claim 26, wherein the antibody is
labeled with a detectable marker.

37. (Previously presented) The method of claim 26, wherein the antibody is
labeled with a radioisotope.

38. (Previously presented) The method of claim 26, wherein the antibody is
labeled with radioactive iodine.